

SOIL

A mulch can be applied when soil temperature is adequate and your first significant irrigation is applied. The mulch will buffer temperature and soil moisture so make sure the correct conditions are buffered. Do not place the mulch against the trunk of the tree. This may promote conditions for pathogen infestation. (In the event of a dry season, the mulch layer may be laid earlier to preserve winter moisture.)

Soil preparation can be done from October, provided the soil moisture is correct. This moisture status is the same as the “1 to 5” ratings as described in the irrigation section of the September Timely Hints. We can prepare heavier soils at a “3” and sandier soils at a “3 and 4” rating.

Important points before soil preparation can take place:


- The area must be cleared of plants, old roots, old structures, drains, old irrigation lines, ridges, etc.
- Any ash piles that remain following the burning of plant material must be spread out to ensure that the ash is not concentrated in specific areas.
- Any old roads within the area must be ripped to relieve compaction. Sample and ameliorate these roads separately.
- A systemic weed killer can be used to kill weeds.
- Apply ameliorants according to your soil scientist’s recommendations.

Two main implement options exist for deep soil preparation actions: a mixing and ripping implement. Depending on the soil conditions and the crop that is to be established, different techniques are recommended. In general, these actions can be described as follows:

- The **loosening action** is generally recommended as the initial action to loosen the soil before a mixing action is implemented. A ripper is generally used.
- The **mixing action** allows for mixing of the soil horizons to make the soil physically uniform. This helps to incorporate the applied ameliorants to a specific depth and so make the soil chemically uniform as well.

Before soil preparation takes place the following points should always be considered:

- Only cultivate the soil at the correct soil moisture content. This is between field water capacity and the lower plastic limit. In a more practical sense, at the correct soil moisture content a ball of soil can be made in the hand and when it is lightly squeezed with the thumb it will crumble. Sandy soils can be slightly wetter than clayey soils, but after the sample is squeezed, no free water must remain in the hand.
- Ensure the slice width between the actions is 2/3 of the working depth. Applying this principle will ensure that the soil across the total area is prepared. Wider slice widths will result in pockets of soil that have not been loosened and ameliorated.



- If cultivating the land in two directions, these actions must always be at a 30-40° angle to each other, with the final action being in a downhill direction.
- Never work deeper than recommended, and never allow clay to come to the soil surface.
- Always make sure to work to the edges of the block during ameliorant and soil preparation. Failing to do so may cause the crops at the edges of the block to not perform well.
- After soil preparation no wheel traffic must be allowed over the land.

These basic descriptions are for the producer's understanding and do not replace the soil scientist's evaluation or recommendation. Executing the incorrect action could drastically reduce the soil's potential for cultivation of fruit crops.

IRRIGATION

- Keep monitoring soil moisture as described in the September *Timely Hints*.
- Be aware of the phenological stage of the cultivars on your farm. From pip hardening to 3 weeks before harvest is the critical period for cell enlargement. Incorrect irrigation during this period will lead to smaller fruit with sub-optimal internal quality. (The pip hardening period is less sensitive for controlled water stress.)

NUTRITION

- For early cultivars, especially nectarines, potassium (K) needs to be available. The period for highest K demand is the same as the critical irrigation period. This is why K availability is often mistaken as the cause for small fruit, resulting in greater K applications the following year with no significant impact on fruit size. The opposite is also true if K is deficient and irrigation optimal.
- Adequate Nitrogen (N) should now also be available. Select the correct source of N:
 - LAN is great for sand and clay soils. Urea should not be used on sandy soils. Calcium nitrate leaches easily and should only be used in sandy soils if the Ca is needed for improved internal quality and should be applied in smaller quantities more often (2 weekly).
- Convert fertigation to granular conventional fertilizer if soil moisture does not allow for the required irrigation.

SUMMER PRUNING

The principles of Summer Pruning:

One would like to invest in growth where one wants it and convert the photosynthetic energy into fruit enlargement and not into wood production. At the same time as achieving this through summer pruning, one will improve the light management in the tree so that the leaves, especially leaves near the fruit can intercept enough light to photosynthesize to their full potential. Judicial summer pruning is an art and when well executed can make a significant difference to stone fruit production.

Peaches and nectarines

Summer pruning of peaches and nectarines should focus on controlling vigorous water shoots in the tree that we often refer to as “leaky pipes”. They are called leaky pipes because they use up a great deal of the trees energy created by photosynthesis. The main aim of summer pruning in peaches and nectarines is to put as much of this energy into the fruit and not into wood production.

The following is recommended:

- Two weeks before harvest or just before Christmas, whichever comes first, cut back all one year old shoot growth thicker than knitting needle thickness to a 10cm stub. This will stop the shoot growing for a while and the energy can then be allocated to the fruit. It also helps with light management and especially with nectarine’s colour development etc. Where the summer pruning has been well executed, we have seen an improvement in fruit size and the sugar content of the fruit, as the light management is much better.

- One should also keep the bowl of the tree, especially at the base, clean of all new water shoot growth right at the base. To remove most of this growth entirely and higher up one can keep it short by stubbing to 10cm as well.
- In young trees (which you still want to fill the area allocated to them), pinching out the growing tip of the lateral shoots coming off the central or closed vase leader so that the terminal shoot remains dominant is important. This probably has to be done a few times during the summer.

Apricots

Apricots are spur bearers and bear on 2 year old wood. One wants to create as much one-year-old wood as possible that one can convert the next year to bearing units. It is recommended that all apricots be summer pruned two weeks before harvest by heading back all one year old shoots knitting needle thickness and thicker to a 10cm stub. After harvest, these shoots will put on a little bit of regrowth. These shoots can be left until the winter pruning, which for apricots should be when the trees are in blossom.

Plums

There has been much debate about plums' summer pruning and there are probably more than 10 recipes out in the field and many consultants will have their own strategy.

The following is recommended:

- For willowy growers like Pioneer and Laetitia, consider a double summer prune, removing strong upright one year old shoots two weeks before harvest, this is to give the trees a so called haircut by cutting back one year old shoots knitting needle thickness and thicker to a 20cm long stub.
- For spur type bearers such as Sapphire and also varieties that tend to give blind wood: prune two weeks before harvest by stubbing all knitting needle thick and thicker one year old shoots to 20cm in length.

All Plums: For all plums that are showing excessive vigour, as soon as possible after harvest, come through and remove those very strong pencil thick and thicker one year old shoots with a 2cm stub ("tappie") cut. Especially if these shoots are growing after harvest, they should be removed. One does run the risk of stimulating shoot growth but in my experience, if you make the stub cut short enough, one normally does not get re-growth and then the photosynthetic energy goes into building up reserves for next year and not into the tree producing more wood.

PEST AND DISEASE CONTROL

- The latest flowering cultivars of all stone fruits may be progressing through the phenological stages of development as depicted in the September “Timely Hints”. Please refer to that document to cover the pest and disease strategies that are to be applied, given those phenological stages of development.
- As the fruit approach maturity, one must be very aware of complying with the registered required withholding periods to prevent chemical residues being detected on the fruit destined for both the export and domestic markets. It is imperative that one consults closely with your exporter to determine what additional MRL restrictions may apply to your fruit for their marketing programmes. Indoxacarb and Thiacloprid are no longer tolerated in the international markets.
- In addition to the powdery mildew, thrips, OFM, FCM and fruit weevil, one must also bear the following in mind:
 - **Brown Rust, Freckle, Gum spot – All Stone Fruit.** Spray 150g/hl Dithane at 75% petal drop and repeat 14 daily for 2-3 sprays, if needed (wet conditions). Safety window = 63 days on peaches, 42 days on apricots and nectarines and 35 days on plums. Be aware of possible discerning market restrictions on Mancozeb usage. Alternatively, for brown rust, spray 14 daily 300g/hl wettable sulphur alternated 14 daily with 200g/hl Captab to control gum spot and freckle. 35 day Safety window applies to Captab and a suggested 35 day window for sulphur to avoid visible spray residues as there is no safety window applicable for sulphur.
 - FCM Monitoring – Plums, peaches and nectarines. This needs to be undertaken from petal drop onwards, for all orchards being produced for EU/UK export. Need 1 trap per orchard to a maximum size of 2 Ha. Separate traps for official monitoring, must be used for FCM and FF. In terms of the industry’s stance on FCM in plums, here too, traps need to be installed to monitor as per the peaches and nectarines.
 - **The FCM Virus Cryptex/Gratham is registered on all stone fruit.** This should be applied 10-14 days after a peak moth catch in the traps, at a rate of 60ml/Ha and be repeated 14 daily thereafter at 35ml/Ha, until the moth infestation is under control. These virus applications should be applied in the later afternoon or evening (owing to their UV sensitivity), to assist in the viral load to suppress the FCM activity. All the FCM remedies, including the virus, are very Ph sensitive, and need to be maintained in the 5-8 range, for optimum efficacy.
 - Broadband or Eco Bb(Beauveria bassiana) is a fungal contact insecticide which must be sprayed on its own. Must add a wetter-sticker like Biodew or Villa 51 @ 6ml/hl to the spray solution. A series of 3 sprays should be applied. If this is done, will suppress red spider mite.
 - Marksman, despite the 7 day safety window, needs to be positioned early in the control programme as the FCM eggs need to be laid on top of the spray for best efficacy.

The table below gives the current list of products registered for FCM control.

<u>Product Rate/hl</u>	<u>Safety (Days)</u>	<u>Number of Sprays</u>	<u>Peaches</u>	<u>Nectarines</u>	<u>Plums</u>	<u>Apricots</u>
Altacor 10g OR Coragen 17.5ml	14	2	√	√	√	
Akito 12,5ml	14	2	√	√	√	
Cypermethrin 10ml	14	2	√	√	√	
Exirel 50ml	7	2	√	√	√	
Marskman 60ml	7	2			√	√
Ampligo 350ml/Ha	14	3			√	
Delegate 20g	7	4 Including Tracer	√	√	√	√
Warlock 100ml	21	4 'Cots 2	√	√	√	√
Broadband 50ml – Min of 1L/Ha OR Eco Bb 100g	0	Minimum of 3	√	√	√	√

- **Pernicious Scale – Peaches, Nectarines and Plums.** This should basically be controlled in the dormant winter period, when there is very little impact on the predators as well as getting better cover onto the leafless tree, where the scale is situated.
- **Late Scale - All Stone Fruit.** Spray Movento @ 40ml/hl from 14 days after 100% petal drop onwards, at first crawler movement for scale. Repeat 4 weeks later if needed. Safety window = 14 days local and export.
- **Fruit Fly – All Stone Fruit.** Hang 1 trap/2 Ha to monitor fruit fly. These traps must be installed from petal drop onwards in all blocks to monitor the fruit fly activity.
- **Fruit Fly – All Stone Fruit.** Monthly baiting should be done throughout the winter (June, July and August), irrespective of trap catches. In May and September, fortnightly baiting should be undertaken. For the period, October till April, weekly baiting needs to be complied with, early in the morning. “Hotspots” on the farms (home gardens), should have the intensity of baiting doubled, throughout the year. When baiting, it is critical to apply 1L/Ha of attractant per bait application. Hymelure OR Loklure at a 2L/hl solution rate, mixed with 175ml/hl Mercaptothion, applied at 50L/Ha of solution, applies the 1L/Ha. The ph of the water solution should be buffered to 4.5 – 5.5 for optimum efficacy of Mercaptothion. When one applies GF120 closer to harvest, again the rate is 1L/Ha of

GF120 in 10 to 20L of spray solution/Ha. The lower the volume of water, the better it works. Alternate rows need to be sprayed, effectively wetting 1 side of every tree when baiting. The bait solution needs to be aimed into the top third of the trees, wetting the underside of the leaves.

- **Fruit Fly Damage Assessment – All Stone Fruit.** In terms of Hortgro’s FF Management System for export, a comprehensive damage assessment of inspecting 10 fruit per tree on the marked 25 trees/2Ha block must be undertaken. This must be done every 14 days, from 6 weeks before expected harvest. The final pre-harvest damage assessment should be done as close to harvest as possible (within a week of harvest).
- **False Codling Moth Damage Assessment – Peaches and nectarines.** In terms of Hortgro’s P/N FMS Management System for export, a comprehensive damage assessment of inspecting 10 fruit per tree on the marked 25 trees/2Ha block must be undertaken. This must be done every 14 days, from 6 weeks before expected harvest. The final pre-harvest damage assessment should be done as close to harvest as possible (within a week of harvest).
- **Postharvest decay – All Stone Fruit.** Apply 80ml/hl Indar, 21 days before expected harvest to cover for Monilinia (Brown rot). Indar has a 1 day export and local safety window.
Apply 2L/Ha Protector, 1 day before harvest, covering for Botrytis and Monilinia. A 1 day safety window applies to both export and local. In the event of the weather being conducive to decay, one could consider rather applying 2L/Ha Tutor, 1 day before harvest, covering for Botrytis, Monilinia and suppression of Rhizopus. Tutor also has a 1 day safety window for both the export and local markets.

This “Stone Fruit Timely Hints” was compiled by Andrew Hacking of Ad Lucem Agricultural Services CC, using agrochemical information obtained from Agri-Intel and Hortgro’s MRL database.

TIMELY HINTS CONTRIBUTORS

<p>Soil, Irrigation, and Nutrition Mico Stander Soil scientist Agrimotion 021 851 1051</p>	<p>Thinning and Newly planted trees Peter Dall Technical consultant Peter Dall Consultancy 028 272 9671</p>	<p>Pest and Disease control Andrew Hacking Technical consultant Ad Lucem Agricultural Services 021 8555 674</p>
---	--	--

DISCLAIMER

This document is issued by HORTGRO (Pty) Ltd (Reg no. 2005/010343/07). Whilst all care has been taken in the provision of the information in this document, this information is provided without liability to us, our affiliates or any officers or employees of ours. The information expressed in this document is done in good faith and is not intended to constitute any form of advice, including but not limited to horticultural practises, such as the administration, management, production and export of products. Some of the information may have been provided or sourced from third parties and we do not in any way guarantee its accuracy or correctness. At all times we will endeavour to ensure that information obtained from any third party is accurate and reliable. HORTGRO and its affiliates disclaims and assumes no liability for any loss or damage (direct, indirect or consequential) that may be suffered from using or relying on the information contained herein. Investment in agriculture and research may cause exposure to certain risks, including market risk, and financial losses, therefore it may not be suitable for all clients. Please contact a technical advisor for a personal analysis prior to making any orchard practice decisions. HORTGRO and its affiliates disclaims and assumes no liability for any loss or damage (direct, indirect or consequential) that may be suffered from using or relying on the information contained herein without seeking professional advice.

