















- Applying herbicides before green tip so that the mites are not forced into the trees when there is foliage for them to feed on.
- Maintaining an organic layer on the orchard floor for the survival of predators during winter by encouraging plants on the orchard floor.

## REFERENCES

- BELCZEWSKI, R. & HARMSSEN, R., 2000. The effect of non-pathogenic fungi on life-history traits of urticae (Acari: Tetranychidae). *Experimental and Applied Acarology* 24, 257–270.
- BROWN, L. & PRINGLE, K.L. 2006. Monitoring system for pests on pome fruit. *Deciduous Pamphlet Series*. DFPT Research, 3pp.
- MATTSON, W.J. & HAACK, R.A., 1987. The role of drought in outbreaks of plant-eating insects. *Bioscience* 37, 110–118.
- PRINGLE, K.L., 2001. Biological control of tetranychid mites in South African Apple orchards. *Acarology: Proceedings of the 10<sup>th</sup> International Congress*. RB Haliday, DE Walter, HC Proctor, RA Norton and MJ Colloff (Editors). CSIRO Publishing, 429-435.
- PRINGLE, K.L. & HEUNIS, J.M., 2006. Biological control of phytophagous mites in apple orchards in the Elgin area of, South Africa using the predatory mite, *Neoseiulus californicus* (McGregor) (Mesostigmata: Phytoseiidae): a benefit–cost analysis. *African Entomology* 14: 113-121.
- PRINGLE, K.L. & HEUNIS, J.M., 2017. Mites with special reference to pears: General considerations. *South African Fruit Journal Dec16/Jan17*: 58-59.
- PRINGLE, K.L. & HEUNIS, J.M., 2017. Mites with special reference to pears: Results from Ceres. *South African Fruit Journal Dec16/Jan17*: 60-63.