



Pruning of peach and nectarine trees



Pruning videos:

- Peach/nectarine trees: <https://youtu.be/CXkqZ03pXZw>
- Apricot trees: <https://youtu.be/SniO4e2XloU>
- Plum trees: <https://youtu.be/Tx4SSAUqTMk>

For all Deciduous Fruit Trees, in actual fact all fruit trees, light management is critical. It is important to remember the saying, we don't farm fruit, we farm light to produce fruit.

The principle of peach and nectarine pruning is that peaches and nectarines crop on one year old wood and one needs to achieve good quality bearing units (one year old shoots) as near to the structural limb as possible:

- No matter what training system one is using, one must identify the structural limb and keep this on a single plane, that is, remove all forks off the structural limbs, be it a central leader tree, a two leader tree or a four leader vase tree. Identify the structural limb and prune leaving one year old knitting needle thick and slightly thicker one year old wood as near to the structural limb as possible.

in length. The nearer these one year old shoots that one is cutting back to is to the structural limb, the better, so the aim must be to have as little two year old and older wood besides the structural wood in the tree.

- If one concentrates on removing the forks, one normally sorts out most of the light problems.
- Remove matchstick thickness and thinner one year old shoots. These thin little bearers just do not have enough reserves and leaves to produce good quality fruit.
- Do not cut back into one year old wood, especially in winter as these cuts normally create dieback.
- Upright, up to 80cm in length one year old shoots are fine as they will bear fruit and then with the weight of the fruit, the shoots will bend down and no longer be vertical.

The best way of checking if one is doing enough pruning or too much pruning, is to calculate the number of fruit a tree can produce per centimetre trunk circumference. Some of the early varieties, one cannot load more than 6 fruit per centimetre trunk circumference but for some of the later highly precocious varieties, one can go up to 14 fruit per centimetre trunk circumference.

It is good to find out from your Technical Advisor what would be the ideal crop load that these trees could carry given the variety, tree density, training system and tree age. One counts all one year old shoots longer than 20cm in length as a bearing unit and works on an average of 1.5 fruit per bearing unit. So for example, if one has a variety that can produce 6 fruit per centimetre trunk circumference given the training system age etc., and the trunk circumference is 30cm then this tree is capable of producing 180 fruit of a desired size per tree. In order to get 180 fruit one needs 120 bearers per tree (1.5 fruit per bearer).

The author likes pruning peaches in early autumn so that the remaining wood can be exposed to direct sunlight and mature properly before going into winter rest. **See picture below** of good quality one year old bearing wood that one is aiming to leave on the tree to carry the next year's crop.



Disinfection of pruning equipment after pruning of each tree is important, one does not want to spread bacterial or other diseases with the pruning shear. You should disinfect with a recommended disinfectant strictly adhering to the label instructions.

With Stone Fruit, the author prefers to paint pruning wounds bigger than a R5 coin, with a suitable tree sealer. The author does not believe it is necessary to paint apple and pear wounds but likes to do this with stone fruit. Never prune on a rainy day as one can spread disease very easily in rainy conditions.

The author has no problems with putting pruning cuttings on the "bankie", either by chipping them with a pruning cutter/slasher that then places chips on the "bankie" or placing them neatly on the "bankie" by hand. The bigger branches can be cut up fine to put on the "bankie". It is a good way of building up the organic material on the "bankie"

Post-harvest and autumn chores

A number of production inputs need to be attended to in the post-harvest/Autumn period. It is good to go through the check list to make sure that they are all actioned:

- Drainage: Make sure all drains are open, unblocked and ready for the winter rains
- Support Trellis: Repair and wires tightened etc.
- Weed Control: Attended to, especially perennial weeds; perennial grasses such as Kikuyu, Fynkweek etc., these are easy to control in Autumn with a good systemic herbicide when the sap flows to the roots.
- Study Cull Analysis Results to determine what the pest and disease problems were in the previous season and start defining a strategy for the coming season. Many pest and diseases are controllable in Autumn or Winter.
- Make sure that all trees are tied firmly to the support trellis, you do not want trees rocking in the ground.
- Tie leaders firmly to support trellis.
- Analyse yields of the previous season and determine which factors contributed to lower than expected yields and how many of those factors can be addressed, and attended to, during the coming Autumn, Winter and Spring.
- For example; cross pollination: Was there adequate cross pollination and did you have the right pollinizers? For plums this is critical. As discussed previously, two pollinizers are better than one. You can determine whether it is necessary to graft in additional cross pollinators in July/August and start making plans to action.
- Nematode Samples: Take Nematode samples and submit for analysis, to determine if this is one of the limiting factors in the orchards.
- Obtain a post-harvest fertilizer and foliar nutritional spray programme from your plant nutrition consultant

Conclusion

In conclusion, post-harvest/autumn is the time that one starts planning for the new season and try to rectify the short comings of this past season. A well

planned strategy for this coming season is invaluable and this should be undertaken in the post-harvest/Autumn period.

TIMELY HINTS contributors:

Soil, Irrigation, Nutrition Mico Stander Soil scientist Agrimotion 021 851 1051	Harvesting tips/Pruning Peter Dall Technical consultant Peter Dall Consultancy 028 272 9671	Pest and Disease control Andrew Hacking Technical consultant Ad Lucem Agricultural Services 021 880 1905
--	--	---

*Disclaimer

This document is issued by HORTGRO (Pty) Ltd (Reg no. 2005/010343/07). Whilst all care has been taken in the provision of the information in this document, this information is provided without liability to us, our affiliates or any officers or employees of ours. The information expressed in this document is done in good faith and is not intended to constitute any form of advice, including but not limited to horticultural practises, such as the administration, management, production and export of products. Some of the information may have been provided or sourced from third parties and we do not in any way guarantee its accuracy or correctness. At all times we will endeavour to ensure that information obtained from any third party is accurate and reliable. HORTGRO and its affiliates disclaims and assumes no liability for any loss or damage (direct, indirect or consequential) that may be suffered from using or relying on the information contained herein.

Investment in agriculture and research may cause exposure to certain risks, including market risk, and financial losses, therefore it may not be suitable for all clients. Please contact a technical advisor for a personal analysis prior to making any orchard practice decisions.

HORTGRO and its affiliates disclaims and assumes no liability for any loss or damage (direct, indirect or consequential) that may be suffered from using or relying on the information contained herein without seeking professional advice.