

# HORTGRO

Growing Fruit IQ

*home of the south african stone fruit producer*



Stone

# TIMELY HINTS

**AUGUST 2018**

STONE FRUIT PRODUCTION

## SOIL

### General maintenance on your farm

- Clean drains of any debris or roots to ensure the drains are working optimally. Wet soils take longer to warm up (>13°C) and this will affect the bud break phase of trees in winter rainfall areas.
- Maintain roads and orchard floors in this period to facilitate tractor and spray cart movement.
- Maintain ridges to allow for an even surface. This will prevent puddling or run-off of irrigation water. Start the re-establishment process as soon as the final harvest is off the orchard in order to allow for the maximum time before planting.

### Last application of lime

- Lime should not be applied much later than the beginning of August.
- Lime has a very low solubility (approximately 2%) and needs ample time (May to September) and water (rain in winter rainfall areas and irrigation in summer rainfall areas) in order to react in the soil and rectify pH within the root zone.

Applying too much lime or applying lime too late may increase adverse interactions between elements. Elements that are negatively affected by very high Calcium (Ca) concentration and pH are Zinc (Zn), Boron (B), Manganese (Mn), Magnesium (Mg), Iron (Fe), Potassium (K), Phosphorus (P). Thus stick to your advisor's recommendations.

## IRRIGATION

### Install irrigation for new developments

- If you have not done so already, install new irrigation systems for new developments.
- Do not wait until it is time to plant.

## Ensure irrigation probes are appropriately placed and correctly installed

- Evaluate each soil moisture probe with regards to:
  - Data quality in the previous season: do you see clear trends?
  - Placement relative to the tree (in root zone?) and emitter (in wetted zone?)
  - Is the probe still in working condition?
- Decide whether any probes need to be repaired, moved, or reinstalled.
- Install any new soil moisture probes. Probes need to be representative of:
  - The largest soil body in the orchard.
  - Primary cultivar in the orchard.
  - Typical trees in the orchard (not a small sickly area or a small vigorous area)

Probes are a vital part of effective water management and it is recommended that you familiarise yourself with the hardware and software and costs (capital and time) of various offerings. BUT please make sure that you are informed when making your decision to use, or not use, soil moisture probes.

## Prepare irrigation system

- Do as much maintenance on the system as possible before the onset of the irrigation season as to avoid mishaps in critical periods.

## NUTRITION

### Order your fertiliser

- Do a stocktake of available fertiliser. Clean the storeroom. Your new fertiliser will arrive soon.
- Order all fertilisers and foliar sprays if you have not done so already.

## REST BREAKING OF PLUMS

This is one of the most critical actions on any stone fruit farm. The aim must be to get the tree to break rest uniformly and as condensed as possible. In South Africa, we never have enough winter chilling for most of our pome fruit varieties and at least half of the stone fruit varieties, so rest breaking is critical.

- All peaches, nectarines, apricots, and plums should be divided into high chilling, medium chilling, or low chilling requirement cultivars and the choice of product must be made according to the chilling requirement of that specific cultivar. You should also take the number of cold units recorded for the year and the delayed foliation history of the orchard [into account].
  - High chilling requirement cultivars such as Gaviota and Southern Belle:
    - ❖ Dormex 0.5 Litres per 100 Litres Water plus
    - ❖ Bud break (or similar product) 2-4 Litres per 100 Litres Water.
  - Medium Chilling Requirement Cultivars such as Songold and Laetitia:
    - ❖ 4 Litres Bud Break (or similar product) per 100 Litres water, or Dormex 0.5 Litres plus 2 Litres Bud Break per 100 Litres water.
  - Low Chilling Requirement Cultivars:
  - 3 Litres Bud Break per 100 Litres water.

On some of the Low Chilling Requirement Cultivars, no rest breaking product is required. This is only applicable in areas with sufficient chill units to satisfy the requirements of cultivars.

- The following should be noted:
  - Rest breaking products to be sprayed at 70-90% of high volume requirement (minimum of 1000 Litres per hectare). Droplets 150 to 250 Micron will give best results.
  - It is important that you should experience some kind of bud swell before spraying but you do not want to spray later than the bud break stage.
  - With Dormex the pH of Spray Water is important, must be below 7.0 pH.
  - Damage is sometimes experienced, to avoid damage, it is essential to get thorough continuous agitation of the spray mixture in the tank.
  - Damage can also occur under the following conditions:
    - ❖ Stress Conditions caused by too wet or too dry soil or young unripe wood which normally occurs where growth continued until late Autumn/Winter.
    - ❖ When applying oil, best results are achieved if one has two to three days of warm weather after the oil spray so that the buds can sweat a bit under the film of oil.

**If one can ensure good rest breaking, this will make farming for the rest of the year a lot easier, especially for thinning and harvesting. The more condensed and uniform the bloom, the better.**

There are other rest breaking products but in my experience, Dormex and Oil provide the best results. After the oil, if one wants to give the trees a boost, you can spray a week later with:

- 1 to 1,5 kg of Potassium Nitrate plus
- 6 ml Biowet per 100 Litres water.
- Very good results have been achieved where one sprays peaches and apricots with rest breaking sprays, normally oil at between 2 to 4 Litres per 100 Litres water is sufficient for most peaches, nectarines, and apricots.

## POLLINATION

All peaches and nectarines are self-fertile and do not require cross-pollinators. Most apricots are self-fertile and don't need pollinators but we have seen the advantage of placing bees into apricot orchards to distribute the pollen in the flower. We have definitely seen higher yields where bees have been put into apricot orchards.

For 80% of plum varieties, cross-pollination is critical. The industry norm is to have at least 11% of a cross pollinator variety in an orchard, often 20% is more desirable and having two cross pollinating varieties in an orchard is also very advantageous as one year the one may be a little bit out of sync with the main variety and the other year the other variety is more in sync. For varieties that are difficult to pollinate like Songold, at least 25% cross-pollinators in the orchard are recommended.

- Cross-pollination is the biggest reason for not achieving the desired yields in a plum orchard. Where you have adequate cross-pollination and good bee hive activity, you can almost be assured of a good yield. This is one production input that you must not skimp on.
- For plums, you will require 3-6 bee hives per hectare from 20% blossom to 50% petal drop. More and more consultants in the field are saying that for difficult to pollinate varieties, we should be aiming for up to 12 hives per hectare. This seems very high but if it assures a good cross pollination, it is worth it.
- The advantage of rest breaking sprays in an orchard especially at high rates, [is that] it synchronises the pollinators much better with the main variety.
- Bee hives must not be put on the damp ground. They should be placed on crates, motor car tyres, or something that will take them off the damp ground. Also, they should ideally face east to catch the early morning sun. You should ensure that you do not apply chemicals that irritate or kill the bees.
- Local knowledge as to what the best pollinators are for the various varieties is your best source of information to determine which pollinators to use for which varieties.
- Spray practices during pollination

Whether a crop needs honeybees for pollination or not, and whether commercial beehives are hired or the producer takes his chances on natural pollinators, and even if the producer does not want pollination producing unwanted pips in certain crops, most flowers will be visited by both unwanted and beneficial insects - mostly in daytime, and whether you want to believe it or not - honeybees are insects as well!

Honeybees collecting nectar and/or pollen will and/or can be affected by all Chemicals - be it a fungicide, insecticide, herbicide, foliar feed, additive, or even just plain water, sprayed from the ground or from the air or sometimes fed through irrigation water into the root zone, applied both before and/or during flowering, with the effect either immediately or through delayed action, manifesting through disruption and/or repellence and/or poisoning.

And as proper pollination (or the lack thereof) does cost money, it will pay off to adhere to the following:

- Think about, plan and execute all chemical applications on all crops with care;
- Refrain from spraying during bloom if at all possible—all crops;
- Flowering weeds are highly attractive to bees—choose the method and timing of control with care;
- If spraying is needed during bloom, do so in the evening after the bees have stopped foraging;
- If using bees for pollination, plan for it beforehand, and place the hives appropriately;
- Do not spray within 2 to 4 full days of introducing the bees at around 15% bloom;
- Take care with snail pellets during and after rain and irrigation—contaminated pools are a serious risk to bees' drinking water.

## PEST AND DISEASE CONTROL

- **Bacterial diseases & gumspot—All stone fruit.** Spray at **first signs of bud movement** Copper Oxychloride @ 350g/hl. Last copper spray for the season whilst fruit is on trees.
- **Scales and mealy bug—All stone fruit.** At **first signs of bud movement**, apply 2nd (final) 100ml/hl Dursban plus 500 ml/hl mineral oil for scale control. Where rest breaking sprays are applied to apricots and plums, rather apply this treatment with the heavier oil spray, effectively negating the 500ml/hl mineral oil requirement. If no rest breaking sprays are applied, add this spray to the copper.

- **Late Scale and Aphids—Peaches, nectarines and plums.** Spray Closer @ 12ml/hl from 100% petal drop onwards for scale. If needed repeat 6 weeks later. Closer @ 5ml/hl is the aphid dose. Safety window = 14 days.
- **Leaf curl—Peaches and nectarines.** Spray Thiram @ 150g/hl **5 to 7 days after copper** and **repeat every 4 to 7 days** (based on how wet trees are—dew and rain) before and after rain. Last spray 90% Petal Drop.
- **Blossom blight (*Monolinia laxa* brown rot)—All stone fruit.** Spray Chronos @ 27 ml/hl **weekly from 1 week after the copper till full bloom** on apricots and plums. In the case of peaches and nectarines add Chronos to the 30% blossom Thiram spray.
- **Powdery mildew—Peaches, nectarines, and apricots.** From **10% balloon on**, apply 60 ml/hl Nimrod @ 10-14 day intervals till 90% petal drop. Apply wettable sulphur @ 300 g/hl from 90% petal drop onwards at 10-14 day intervals, till 35 days before expected harvest, if required. This wettable sulphur will also control brown rust.
- **Powdery Mildew—Apricots.** From **10% balloon on**, apply 1L/Ha Ortiva @ 10-14 day intervals, applying no more than 2 sequential sprays and 3 sprays in total, till 90% petal drop. From a resistance point of view, one should add 300g/hl wettable sulphur to these Ortiva sprays. Wettable sulphur is registered for brown rust control and as such will assist in controlling mildew. In addition to these sprays, apply wettable sulphur @ 300g/hl from 90% petal drop onwards at 10-14 day intervals, till 35 days before expected harvest, if required.
- **Blossom complex pests (Bollworm, Antestia, Fruit Nibbler, Green Peach Aphid)—Nectarines, Peaches and Plums:** Spray Klartan @ 30 ml/hl @ 10–30% flower (before bees are put into orchards), as and where required. Do not spray after 90% petal drop. Klartan is “safe” for bees.
- **Blossom complex pests (Bollworm, Antestia, Fruit Nibbler)—Apricots.** Spray Delegate @ 12g/hl. This dose will control thrips. A 20g/hl dose is registered for FCM. Safety window = 7 days. After 75% petal drop Lamda-cyhalothrin may be applied @ 10ml/hl for Bollworm or 20ml/hl for weevil control.
- **Blossom Complex Pests (Green Peach Aphid)—Apricots.** Spray Aphox @ 50g/hl when pest first noticed.
- **Thrips—Nectarines and plums.** Apply 15 ml/hl Tracer @ 10% balloon and repeat 7-10 days later if needed. If bollworm is a problem, the Tracer can be sprayed at 20 ml/hl to cover for both thrips and bollworm. Safety window = 21 days on peaches and plums, 7 days on nectarines.

- **Oriental fruit moth (OFM)—Peaches and nectarines.** Where required, hang traps (1 trap/2 Ha) early August at the latest.
- **False codling moth (FCM)—All stone fruit.** Where required, hang traps (1 trap/2 Ha) early August at the latest.
- **Brown rust, freckle, gum spot—All stone fruit.** Spray 200g/hl Captab at 75% petal drop and repeat 14 daily for 2-3 sprays, if needed (wet conditions). Safety window = 35 days on stone fruit. **OR**
- **Brown rust, freckle, gum spot—All stone fruit.** Spray 150 g/hl Dithane at 75% petal drop and repeat 14 daily for 2 to 3 sprays, if needed (wet conditions). Safety window = 63 days on peaches, 42 days on apricots and nectarines and 35 days on plums.
- **Fruit weevil (snout beetle)—Nectarines.** Stem bands are the best form of weevil control and should be placed around tree trunks before the trees start to bloom. You need to ensure that there are no “ladders” of pruning shoots or weeds, allowing weevils to access the trees above the stem bands. For chemical control, spray 40 ml/hl Steward once weevils are caught in the monitor bands or @ 75% petal drop. Safety window = 28 days.
- **Fruit weevil (snout beetle)—Apricots.** Apply 20ml/hl Lamda-cyhalothrin when damage occurs as monitored above from petal drop on. Safety period = 28 days.
- **Crown gall—All stone fruit.** All new trees’ roots should be dipped in RAS 84, before planting. One packet treats 50 trees. *Agrobacterium radiobacter*, a biological treatment is only preventative and not curative.

## TIMELY HINTS CONTRIBUTORS

Soil, Irrigation, and Nutrition	Thinning and Newly planted trees	Pest and Disease control
Mico Stander Soil scientist Agrimotion 021 851 1051	Peter Dall Technical consultant Peter Dall Consultancy 028 272 9671	Andrew Hacking Technical consultant Ad Lucem Agricultural Services 021 880 1905

### DISCLAIMER

This document is issued by HORTGRO (Pty) Ltd (Reg no. 2005/010343/07). Whilst all care has been taken in the provision of the information in this document, this information is provided without liability to us, our affiliates or any officers or employees of ours. The information expressed in this document is done in good faith and is not intended to constitute any form of advice, including but not limited to horticultural practices, such as the administration, management, production and export of products. Some of the information may have been provided or sourced from third parties and we do not in any way guarantee its accuracy or correctness. At all times we will endeavor to ensure that information obtained from any third party is accurate and reliable. HORTGRO and its affiliates disclaims and assumes no liability for any loss or damage (direct, indirect or consequential) that may be suffered from using or relying on the information contained herein. Investment in agriculture and research may cause exposure to certain risks, including market risk, and financial losses, therefore it may not be suitable for all clients. Please contact a technical advisor for a personal analysis prior to making any orchard practice decisions. HORTGRO and its affiliates disclaims and assumes no liability for any loss or damage (direct, indirect or consequential) that may be suffered from using or relying on the information contained herein without seeking professional advice.